

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/281480897>

Evaluation of Apoptosis Induced by Celecoxib Loaded Liposomes in Isolated Rat Hepatocytes

Article in *Jundishapur Journal of Natural Pharmaceutical Products* · September 2015

DOI: 10.17795/jjnpp-25421

CITATION

1

READS

120

7 authors, including:



Mojtaba Kalantar

Ahvaz Jundishapur University of Medical Sciences

3 PUBLICATIONS 4 CITATIONS

SEE PROFILE



Mohsen Rezaei

Tarbiat Modares University

108 PUBLICATIONS 1,239 CITATIONS

SEE PROFILE



Eskandar Moghimipour

Ahvaz Jundishapur University of Medical Sciences

219 PUBLICATIONS 2,368 CITATIONS

SEE PROFILE



Neda Bavarsad

Ahvaz Jundishapur University of Medical Sciences

29 PUBLICATIONS 605 CITATIONS

SEE PROFILE

Evaluation of Apoptosis Induced by Celecoxib Loaded Liposomes in Isolated Rat Hepatocytes

Mojtaba Kalantar¹; Mohsen Rezaei^{1,2,*}; Eskandar Moghimipour³; Neda Bavarsad⁴; Heibatallah Kalantari¹; Golnaz Varnaseri¹; Arash Forouzan⁵

¹Department of Pharmacology and Toxicology, School of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

²Department of Toxicology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, IR Iran

³Department of Pharmaceutics, Faculty of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

⁴Nanotechnology Research Center, School of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

⁵Department of Emergency, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

*Corresponding author: Mohsen Rezaei, Department of Pharmacology and Toxicology School of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran. Tel: +98-9167250548, Fax: +98-6113738381, E-mail: mohsen.rezaei@gmail.com

Received: January 18, 2015; Revised: June 10, 2015; Accepted: July 12, 2015

Background: Apoptosis is an essential process for elimination of damaged and cancerous cells and is also a desired effect for the anti-cancer activity of drugs. Cyclooxygenase-2 (COX-2) inhibitors including celecoxib are a class of drugs with chemopreventive and anti-proliferative properties regardless of their routine anti-inflammatory effects.

Objectives: The objective of this study was to evaluate the effect of celecoxib loaded nano-liposomes on the isolated rat hepatocytes and comparing the results with what were obtained from usual form of this chemopreventive agent.

Materials and Methods: Freshly isolated rat hepatocytes were prepared by two step collagenase perfusion method and, following stabilization in rotary, were exposed to 0, 20, 40 and 100 μ M of celecoxib in nano and usual form. Viability was obtained and apoptosis was determined by modified comet technique.

Results: At high concentrations (40 and 100 μ M), apoptosis was demonstrated and it was clearly more prominent in hepatocytes exposed to celecoxib loaded liposomes.

Conclusions: Our results showed that celecoxib loaded liposomes robustly induced apoptosis, an action that can potentially make it more relevant than the usual form for chemopreventive strategies and also may raise concerns about its toxicity.

Keywords: Apoptosis; Celecoxib; Liposomes; Hepatocyte; Comet Assay; Cyclooxygenase-2

1. Background

Nanotechnology involves imaging, measuring, modeling and manipulating material at nanoscale (1). Nanoscale materials have unique physicochemical properties, such as high reactivity, ultra-small size, and larger ratio of surface area to mass, which are different from bulk materials (in microscale) of the same composition. These properties can be useful to decrease several of the limitations found in nanomedicine (2, 3). Nanomedicine has been defined as the application of nanotechnology to medicine. The difference between nano-drugs and conventional drugs is that nano-drugs are entirely based on small molecule chemistry. By manufacturing drugs in nanoscale, is possible to modify their pharmacokinetics and to improve the efficacy and safety of the therapy (4). Nanosized drugs, because of the large surface area to mass ratio, increased the effectiveness of drugs. The nanoparticles are good carriers for poorly soluble compounds (5). Current researches show that the bulk materials are more reactive and have more toxic effects than smaller particles (6, 7). Nanoparticles are not competently scavenged by macrophages, and therefore their blood circulation time is increased. Prolonged bio-

availability extends the duration of controlled system drug delivery or improves the characteristics for nanoparticles to reach target sites, by extravasation (8). Nanotoxicology is rising as an important sub-discipline of nanotechnology. Nanotoxicology deals with the interaction of nanostructures with biological systems and shows the relationship between the physical and chemical properties of nanostructures and their toxic consequences (3, 4). Because of the importance of nanotechnology and its wide use, in the pharmaceutical industry, and the damage caused by the toxicity of nano-drugs, they are receiving increased attention. The coxibs are selective inhibitors of cyclooxygenase-2 (COX-2) (9). The COX₂ inhibitors are used as a treatment for inflammatory diseases, menstrual pain, headache, and patent ductus arteriosus, in premature infants.

Traditional non-steroidal anti-inflammatory drugs (NSAIDs) are prescribed for the treatment of mild to moderate pain, especially pain caused by inflammation. In diseases, such as rheumatoid arthritis and gout, non-selective NSAIDs and COX-2 inhibitors reduced the formation of prolyps in susceptible patients. Also, they reduce the for-

mation, growth and metastasis, in vitro, and also reduced the number of intestinal tumors, in patients with adenomatous polyps (10-12). Apparently, by inhibiting the prostaglandin production and apoptosis induction, NSAIDs exert their anti-tumor effects (13). The NSAIDs reduce the incidence of carcinogen-induced colon tumors (14). Celecoxib is a sulfonamide NSAID and selective COX-2 inhibitor (1). Celecoxib enhances the induction of apoptosis in human pancreatic cancer cells, through a mechanism depending on the prostaglandin E 2 (PGE2) and COX-2 pathways (15), with improvement of colorectal polyps in familial adenomatous polyposis (13). As a chemoprevention of colorectal cancer, it induces apoptosis via calcium ion homeostasis (16). Celecoxib induces apoptosis in cervical tumor cells and colorectal cancer, by blocking the activation of Akt in human prostate cancer cells (17, 18). Target molecules for NSAIDs that participate at signaling pathways, include: 15-lipoxygenase-1, peroxisome proliferator-activated receptors (PPARs), P₂₁ ras signaling and AKT/PKB kinase (19-21). Cerium oxide nanoparticles induced apoptosis and autophagy in human peripheral blood monocytes (22). Doxorubicin nanoparticles have the potential to serve as a beneficial healing approach to overcome the chemoresistance of adriamycin-resistant breast cancer (23).

2. Objectives

The objective of this study was to evaluate the effect of celecoxib loaded nano-liposomes on the isolated rat hepatocytes and comparing the results with what were obtained from usual form of this chemopreventive agent.

3. Materials and Methods

3.1. Chemicals

Triton X-100, dimethyl sulfoxide (DMSO), ethidium bromide, NaCl, KCl, MgSO₄, KH₂PO₄, Na₂HPO₄, NaHCO₃, HEPES buffer, BSA, collagenase, Na₂Cr₂O₇, NaOH and disodium ethylenediaminetetraacetic acid (EDTA), ethylene glycol tetraacetic acid (EGTA), KH₂PO₄, tris base, CaCl₂, sodium lauryl sarcosinate (C₁₅H₂₈NO₃Na) and, Na₂SO₄, low melting point agar, normal melting point agar were obtained from Sigma (Darmstadt, Germany), celecoxib loaded liposomes were prepared in the School of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran. Liposomes were prepared by thin film method. Briefly, soya lecithin and cholesterol were dissolved in chloroform-methanol (1:1) and 5 mg celecoxib was added in the solution, then the mixture was evaporated in a rotary evaporator (24). Hepatocytes extraction solution: Hanks × 10 buffers: composed of KH₂PO₄ (0.3 g), NaCl (40 g), Na₂ HPO₄ (0.3 g), KCl (2 g), MgSO₄, 7H₂O (1 g) and dH₂O q.s. to 500 mL. Hanks I buffer: composed of dH₂O q.s. to 500 mL, HEPES (1.5 g), NaHCO₃ (1.05 g), and Hanks × 10 (50 mL). Krebs solution: composed of MgSO₄ (0.295 g), NaCl (6.95 g), KCl (0.355 g), (0.287 g), KH₂PO₄ (0.16 g), CaCl₂ NaHCO₃ (2.1 g), and dH₂O q.s. to 1000

mL. Perfusion buffer: composed of BSA (1.3 g), EGTA (0.039 g), and Hanks buffer q.s. to 200 mL. Collagenase solution: composed of CaCl₂ (0.03 g), collagenase (0.065 g), and Hanks I buffer q.s. to 100 mL. Washing solution: composed of Krebs buffer q.s. to 200 mL, BSA (2 g), and HEPES (0.06 g). Incubation solution: composed of HEPES (0.75 g) and Krebs buffer q.s. to 250 mL. Neutral comet assay solution include: 1) phosphate solution: composed of NaCl (0.8 mg), KCl (0.02 g), Na₂HPO₄·2H₂O (0.216 g), KH₂PO₄ (0.02 g), and dH₂O q.s. to 100 mL; 2) lysis buffer: composed of NaCl (14.61 g), Na₂EDTA (3.72 g), Tris base (0.121 g), Sodium lauryl sarcosinate 1% (1 g), NaOH (0.9 g), DMSO 10% (10 mL), Triton X-100 1% (1 mL), and dH₂O q.s. to 100 mL. Electrophoresis solution: composed of tris/borate/EDTA (TBE) solution and dH₂O, at a ratio 1:5 (V/V).

3.2. Animals

Male Wistar rats, weighing 230 - 270 g were obtained from Jundishapur University of Medical Sciences, Ahvaz, Iran, and housed in ventilated plastic cages and at temperature of 20 - 24 °C, 12 hour night - 12 hour day light cycle, with controlled humidity and were fed ad libitum.

3.3. Isolated Rat Hepatocytes

Animals were anesthetized by intraperitoneal injection of ketamine 90 mg/kg and xylazine 10 mg/kg. After cannulation of portal vein, liver was perfused, with perfusion solution for 10 minutes and then, with collagenase solution, for 12 minute. Cellular suspensions were prepared and filtered and finally were centrifuged at 1500 rpm, for 3 minutes. Hepatocyte viability was assessed microscopically, by the trypan blue test. The hepatocytes used were at least 85–95% viable. Isolated hepatocytes (10 mL, 10⁶ cells/mL) were placed in a bioreactor bath, under an atmosphere of 95% O₂, 10% N₂, and 5% CO₂, in a 37°C water bath, for 30 minutes.

3.4. Drug Treatment

Three doses of celecoxib loaded liposomes and celecoxib including, 20, 40, and 100 μM (test groups) were prepared and hepatocytes were exposed to different doses of celecoxib loaded liposomes and celecoxib for 60 minute. Incubation buffer was used, as negative, respectively. To determine whether the use of liposome nanoparticles would affect hepatocytes they were exposed to concentrations of 20, 40, and 100 μM of liposomes without the drug, and the results were compared with negative controls.

3.5. Neutral Comet Assay

A volume of 1 mL of cell suspension was mixed with 10 mL of low melting point agarose and 100 μL of this suspension were transferred to a microscope slide, covered with a layer of normal-melting-point agarose. After putting a coverslip on the slides, they were put in the freezer for 10 minute, to allow the agarose to solidify. Then, the coverslip

was carefully removed. Slides were immersed in cold lysing buffer for 1 hour, in the dark. Then, slides were placed side-by-side in a horizontal gel electrophoresis tank, which was filled with fresh electrophoresis buffer. Electrophoresis was conducted for 25 minutes at 25 V. After all these steps, the slides were neutralized in the neutralization buffer and then stained with 50 μ L of ethidium bromide (20 μ g/mL) for 5 minutes, before being analyzed with a fluorescence microscope (25). A total of 50 individual cells were analyzed by the comet assay. An intact cell resembles an undamaged nucleus, without a tail, whereas an injured cell has the appearance of a comet. The apoptotic cells were classified by scoring them as 2 or 3. Score of 0 and 1 reflected intact cells.

3.6. Statistical Analysis

All data were analyzed using the statistical with SPSS version 18.0 (SPSS Inc., Chicago, IL, USA). The results were expressed as means \pm standard deviation (SD) and analysis performed using the non-parametric X_2 test to compare groups of data.

4. Results

In this study, the neutral comet assay was used for the comparative assessment of the apoptotic induction effect of the usual form of celecoxib with celecoxib loaded liposomes in the rat hepatocytes. Totally, 85.98% of cells were

apoptotic after treatment with 100 μ g/mL celecoxib loaded liposomes (Table 1). From images of the fluorescence microscope, apoptotic cells showed comet like images, with long tail and small head, while intact cells were without any tail.

Celecoxib loaded liposomes induced an apoptotic response in hepatocytes and produced a comet pattern, similar to the usual form of celecoxib. The percentage of normal and apoptotic cells are shown in Table 1. In this study, hepatocytes were initially exposed to 20, 40, and 100 μ M of celecoxib loaded liposomes. As shown in Table 1, and Figure 1 and 2, the highest percentage of apoptotic cells was achieved at 100 μ M. Celecoxib loaded liposomes, in this concentration, caused an 85.98% apoptotic response and apoptotic cells showed no statistically significant differences, compared to usual form of celecoxib ($P < 0.0001$). Also, there were significant differences in apoptosis induction at 20 and 40 μ M dose, between the usual form of celecoxib and celecoxib loaded liposomes ($P < 0.005$). Apoptotic effect of 20 μ M of usual form was evident in 2% of cells. At the lowest concentration of celecoxib loaded liposomes, 20 μ M, the apoptotic effect was evident in $>10\%$ of cells. For a more complete evaluation, hepatocytes were treated with 40 and 100 μ M of celecoxib loaded liposomes. The maximum apoptosis was observed in the presence of 100 μ M dose. This concentration caused a significant increase in apoptotic cells and the percentage of apoptotic cells increased to 85.98%, after treatment ($P < 0.005$).

Table 1. Percentage of Intact and Apoptotic Cells After Exposure to Usual Form of Celecoxib and Celecoxib Nanoparticles^a

Groups	Intact Cell	Apoptotic Cell
Usual form of celecoxib 20 μ M	97.98 \pm 4.14	2 \pm 0
Usual form of celecoxib 40 μ M	95.98 \pm 3.18	3.98 \pm 1.1
Usual form of celecoxib 100 μ M	19.98 \pm 1.72	79.98 \pm 2.67
Celecoxib loaded liposomes 20 μ M	87.32 \pm 4.57	12.66 \pm 2.15
Celecoxib loaded liposomes 40 μ M	85.98 \pm 4.6	13.98 \pm 2.3
Celecoxib loaded liposomes 100 μ M	11.98 \pm 5.56	85.98 \pm 5.03

^a The values are presented as mean \pm SD.




Figure 1. Frequency of Comet Patterns of Hepatocytes After Exposure to Usual Form of Celecoxib and Celecoxib Loaded Liposomes.




Figure 2. Percentage of Apoptotic Cells After Exposure to Usual Form of Celecoxib and Celecoxib Loaded Liposomes.

5. Discussion

Many chronic diseases were linked to the defects in the apoptosis process (28). Coxibs specifically inhibit COX-2 (29). The results of Steinbach et al. showed celecoxib can be employed to improve colorectal polyps in familial adenomatous polyposis (13). The results of Vaish showed that celecoxib and sulindac, as the chemoprevention, caused induction of apoptosis in colorectal carcinomas, via the calcium ion homeostasis (16). Kim et al. (30) showed that caspase-8 and -9 executed the apoptotic effect of celecoxib, in cervical cell line. This necessitates the fas-associated death domain-dependent pathway, in a cell type-specific manner. Also, nuclear factor κ B (NF κ B) may play a key role in celecoxib-induced apoptosis. Piplani et al. (31) studies have shown that, dolastatin 15, a mollusk linear peptide, and celecoxib induced apoptosis via inducible nitric oxide synthase, therefore, revealing the potential of this inflammatory protein, as an antineoplastic target in colon cancer. Hsu et al. (18) data demonstrate that inhibition of Akt activation, in human prostate cancer cells, may play a crucial role in the induction of apoptosis by celecoxib. Investigations showed that several other molecular targets of the NSAIDs, except for COX-2, may exist, such as the signaling pathway that contains, 15-lipoxygenase-1, P21 Ras, PPAR delta, and AKT/PKB Kinase (19, 20, 32). The results of this study are comparable with other researches on cell death effects of celecoxib. The results of this study indicated a dose-dependent increase in apoptosis in rat hepatocytes. Our results showed that the usual form of celecoxib (100 μ M) treatment caused 79.98% cell death in rat hepatocytes. By high concentrations (40 and 100 μ M), apoptosis was demonstrated and it was clearly more prominent in hepatocytes exposed to celecoxib loaded liposomes. Zhou et al. (34) showed that high doses of celecoxib have dose dependently induced apoptosis in the human glioma cell line U251. Li et al. (35) Our results showed that celecoxib loaded liposomes robustly induced apoptosis, which can potentially make it more relevant than the usual form for chemopreventive strategies.

Acknowledgements

Special thanks to the research deputy of Ahvaz Jundishapur university of medical sciences, Ahvaz, Iran, for their financial support.

Authors' Contributions

Study concept and design: Mohsen Rezaei, Haibatolla Kalantari. Analysis and interpretation of data: Mojtaba Kalantar, Golnaz Varnaseri. Drafting of the manuscript: Mojtaba Kalanatar, Arash Frouzan. Liposome preparation and Characterization: Eskandar Moghimipour, Neda Bavarasad. Critical revision of the manuscript for important intellectual content: Mohsen Rezaei .

Funding/Support

This study was supported in part by grant N-85 from the Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

References

1. Ferreira AJ, Cemlyn-Jones J, Robalo Cordeiro C. Nanoparticles, nanotechnology and pulmonary nanotoxicology. *Rev Port Pneumol.* 2013;**19**(1):28–37.
2. Bottini M, Bruckner S, Nika K, Bottini N, Bellucci S, Magrini A, et al. Multi-walled carbon nanotubes induce T lymphocyte apoptosis. *Toxicol Lett.* 2006;**160**(2):121–6.
3. Aillon KL, Xie Y, El-Gendy N, Berkland CJ, Forrest ML. Effects of nanomaterial physicochemical properties on in vivo toxicity. *Adv Drug Deliv Rev.* 2009;**61**(6):457–66.
4. Linkov I, Satterstrom FK, Corey LM. Nanotoxicology and nanomedicine: making hard decisions. *Nanomedicine.* 2008;**4**(2):167–71.
5. Torchilin VP, Lukyanov AN, Gao Z, Papahadjopoulos-Sternberg B. Immunomicelles: targeted pharmaceutical carriers for poorly soluble drugs. *Proc Natl Acad Sci U S A.* 2003;**100**(10):6039–44.
6. El-Ansary A, Al-Daihan S. On the toxicity of therapeutically used nanoparticles: an overview. *J Toxicol.* 2009;**2009**:754810.
7. Chan VS. Nanomedicine: An unresolved regulatory issue. *Regul Toxicol Pharmacol.* 2006;**46**(3):218–24.
8. Borm PJ, Muller-Schulte D. Nanoparticles in drug delivery and environmental exposure: same size, same risks? *Nanomedicine (Lond).* 2006;**1**(2):235–49.
9. Yamakawa N, Suemasu S, Okamoto Y, Tanaka K, Ishihara T, Asano T, et al. Synthesis and biological evaluation of derivatives of 2-{2-fluoro-4-[(2-oxocyclopentyl)methyl]phenyl}propanoic acid: nonsteroidal anti-inflammatory drugs with low gastric ulcerogenic activity. *J Med Chem.* 2012;**55**(11):5143–50.
10. Fischer SM, Lo HH, Gordon GB, Seibert K, Kelloff G, Lubet RA, et al. Chemopreventive activity of celecoxib, a specific cyclooxygenase-2 inhibitor, and indomethacin against ultraviolet light-induced skin carcinogenesis. *Mol Carcinog.* 1999;**25**(4):231–40.
11. Strillacci A, Griffoni C, Lazzarini G, Valerii MC, Di Molfetta S, Rizzello F, et al. Selective cyclooxygenase-2 silencing mediated by engineered E. coli and RNA interference induces anti-tumour effects in human colon cancer cells. *Br J Cancer.* 2010;**103**(7):975–86.
12. Thun MJ, Henley SJ, Patrono C. Nonsteroidal anti-inflammatory drugs as anticancer agents: mechanistic, pharmacologic, and clinical issues. *J Natl Cancer Inst.* 2002;**94**(4):252–66.
13. Steinbach G, Lynch PM, Phillips RK, Wallace MH, Hawk E, Gordon GB, et al. The effect of celecoxib, a cyclooxygenase-2 inhibitor, in familial adenomatous polyposis. *N Engl J Med.* 2000;**342**(26):1946–52.
14. Gaosong W, Jilin Y, Fang D, Shengquan Z, Xingrui L. Celecoxib inhibits proliferation and induces apoptosis via cyclooxygenase-2 pathway in human pancreatic carcinoma cells. *J Huazhong Univ Sci Technol Med Sci.* 2005;**25**(1):42–4.
15. Jendrossek V. Targeting apoptosis pathways by Celecoxib in cancer. *Cancer Lett.* 2013;**332**(2):313–24.
16. Vaish V, Sanyal SN. Role of Sulindac and Celecoxib in chemoprevention of colorectal cancer via intrinsic pathway of apoptosis: exploring NHE-1, intracellular calcium homeostasis and Calpain 9. *Biomed Pharmacother.* 2012;**66**(2):116–30.
17. Piplani H, Vaish V, Sanyal SN. Dolastatin 15, a mollusk linear peptide, and Celecoxib, a selective cyclooxygenase-2 inhibitor, prevent preneoplastic colonic lesions and induce apoptosis through inhibition of the regulatory transcription factor NF-kappaB and an inflammatory protein, iNOS. *Eur J Cancer Prev.* 2012;**21**(6):511–22.
18. Hsu AL. The Cyclooxygenase-2 Inhibitor Celecoxib Induces Apoptosis by Blocking Akt Activation in Human Prostate Cancer Cells Independently of Bcl-2. *J Biol Chem.* 2000;**275**(15):11397–403.
19. He TC, Chan TA, Vogelstein B, Kinzler KW. PPAR δ Is an APC-Regulated Target of Nonsteroidal Anti-Inflammatory Drugs. *Cell.* 1999;**99**(3):335–45.

20. Herrmann C, Block C, Geisen C, Haas K, Weber C, Winde G, et al. Sulindac sulfide inhibits Ras signaling. *Oncogene*. 1998;**17**(14):1769–76.
21. Liu DB, Hu GY, Long GX, Qiu H, Mei Q, Hu GQ. Celecoxib induces apoptosis and cell-cycle arrest in nasopharyngeal carcinoma cell lines via inhibition of STAT3 phosphorylation. *Acta Pharmacol Sin*. 2012;**33**(5):682–90.
22. Hussain S, Al-Nsour F, Rice AB, Marshburn J, Yingling B, Ji Z, et al. Cerium dioxide nanoparticles induce apoptosis and autophagy in human peripheral blood monocytes. *ACS Nano*. 2012;**6**(7):5820–9.
23. Kang KW, Chun MK, Kim O, Subedi RK, Ahn SG, Yoon JH, et al. Doxorubicin-loaded solid lipid nanoparticles to overcome multidrug resistance in cancer therapy. *Nanomedicine*. 2010;**6**(2):210–3.
24. Moghimipour E, Handali S. Utilization of thin film method for preparation of celecoxib loaded liposomes. *Adv Pharm Bull*. 2012;**2**(1):93–8.
25. Azqueta A, Collins AR. The essential comet assay: a comprehensive guide to measuring DNA damage and repair. *Arch Toxicol*. 2013;**87**(6):949–68.
26. Lorenzo Y, Costa S, Collins AR, Azqueta A. The comet assay, DNA damage, DNA repair and cytotoxicity: hedgehogs are not always dead. *Mutagenesis*. 2013;**28**(4):427–32.
27. Heibatullah K, Marzieh P, Arefeh I, Ebrahim M. Genotoxicity determinations of coriander drop and extract of Coriander Sativum cultured fibroblast of rat embryo by comet assay. *Saudi Pharmaceutical Journal*. 2008.
28. Wong V, Szeto Y, Collins AR, Benzie I. The comet assay: a biomonitoring tool for nutraceutical research. *Curr Top Nutraceutical Res*. 2005;**3**(1):1–14.
29. Wood AJ, FitzGerald GA, Patrono C. The coxibs, selective inhibitors of cyclooxygenase-2. *N Engl J Med*. 2001;**345**(6):433–42.
30. Kim SH, Song SH, Kim SG, Chun KS, Lim SY, Na HK, et al. Celecoxib induces apoptosis in cervical cancer cells independent of cyclooxygenase using NF-kappaB as a possible target. *J Cancer Res Clin Oncol*. 2004;**130**(9):551–60.
31. Piplani H, Rana C, Vaish V, Vaiphei K, Sanyal SN. Dolastatin, along with Celecoxib, stimulates apoptosis by a mechanism involving oxidative stress, membrane potential change and PI3-K/AKT pathway down regulation. *Biochim Biophys Acta*. 2013;**1830**(11):5142–56.
32. Shureiqi I, Chen D, Lotan R, Yang P, Newman RA, Fischer SM, et al. 15-Lipoxygenase-1 mediates nonsteroidal anti-inflammatory drug-induced apoptosis independently of cyclooxygenase-2 in colon cancer cells. *Cancer Res*. 2000;**60**(24):6846–50.
33. Butler M. *Animal cell culture and technology*. United Kingdom: Taylor & Francis; 2003.
34. Zhou R, Zhang LZ, Wang RZ. Effect of celecoxib on proliferation, apoptosis, and survivin expression in human glioma cell line U251. *Chin J Cancer*. 2010;**29**(3):294–9.
35. Li WZ, Wang XY, Ding YQ. [Celecoxib induces cell cycle arrest and apoptosis in human tongue squamous carcinoma cell line Tca8113]. *Nan Fang Yi Ke Da Xue Xue Bao*. 2010;**30**(3):486–9.